

Correlation Functions in Holographic Renormalization Group Flows

Wolfgang Mück¹

*Dipartimento di Scienze Fisiche, Università di Napoli “Federico II”
Via Cintia, 80126 Napoli, Italy*

Abstract

We consider the holographic duality for a generic bulk theory of scalars coupled to gravity. By studying the fluctuations around Poincaré invariant backgrounds with non-vanishing scalars, with the scalar and metric boundary conditions considered as being independent, we obtain all one- and two-point functions in the dual renormalization group flows of the boundary field theory. Operator and vev flows are explicitly distinguished by means of the physical condensates. The method is applied to the GPPZ and Coulomb branch flows, and field theoretical expectations are confirmed.

1 Introduction

In the past years, holographic dualities have been used extensively in the study of quantum field theories. According to the holographic principle, the properties of a quantum field theory living on the boundary of a certain bulk space-time are in one-to-one correspondence with the dynamics of a field theory living in that bulk [1,2]. The duality most studied (and very well understood) is the correspondence between conformal field theories and field theories living on anti-de Sitter bulk spaces (AdS/CFT correspondence), *e.g.*, the duality between $\mathcal{N} = 4$ SYM theory and gauged $\mathcal{N} = 8$ super gravity on $AdS_5 \times S^5$ [3–5].

In general, the bulk theories also allow for solutions that describe asymptotically AdS spaces. In the case of $\mathcal{N} = 8$ super gravity, these gravitational backgrounds have been interpreted as the duals of $\mathcal{N} = 4$ SYM theory either perturbed by the insertion of relevant operators or given a non-zero vacuum

¹ E-mail: mueck@na.infn.it

expectation value (vev) of some scalar operator [6–13]. Formulated more generally, asymptotically AdS bulk geometries describe the renormalization group (RG) flows of boundary field theories with conformal UV fixed points [14–22].

Using the recipe of the AdS/CFT correspondence, the correlation functions of operators in boundary field theories with RG flows can be obtained by studying the fluctuations in the bulk theory around the background flow solutions. This procedure has been applied to the operator flow of Girardello, Petrini, Porrati and Zaffaroni (GPPZ) [12] as well as to some of the Coulomb branch vacuum flows of $\mathcal{N} = 4$ SYM theory [11,23]. The correlation functions obtained so far include various two-point functions for scalar operators and the traceless transversal parts of the two-point functions of the energy momentum tensor [24–26,11,27–30], as well as the two-point functions of $\mathcal{R}$ -currents [31]. In [30], also the fermion, vector and gravitino sectors of the super gravity fluctuations are discussed.

In the present paper, we shall consider in detail a generic bulk theory consisting of scalars coupled to gravity. Our improvements with respect to preceding work [28–30] will be two-fold. First, although the fluctuations of the active scalars (scalars having a non-zero background) are coupled to the metric fluctuations, their respective boundary values are prescribed independently, since they are the sources in the boundary field theory. This enables us to obtain, in addition to the scalar two-point function, the complete energy momentum tensor two-point function as well as the mixed scalar–energy momentum tensor two-point function. Second, we are able to distinguish explicitly between the operator and the vev flows. In the former, the boundary field theory is perturbed by inserting a scalar operator into the boundary field theory action with a finite coupling, whereas in the latter there is a change of the vacuum of the field theory. While all one-point functions in the operators flows vanish, the vev flows explicitly exhibit a condensate. These one-point functions are generated by the addition of counter terms necessary to render the generating functional finite.

As a preliminary, it is instructive to consider the foundation of holography, namely the correspondence formula

$$\left\langle e^{-\int \sqrt{\hat{g}} d^d x \mathcal{O}(x) \hat{\phi}(x)} \right\rangle_0 \equiv e^{-S[\hat{\phi}, \hat{g}_{ij}]} . \quad (1)$$

On the right hand side, $S[\hat{\phi}, \hat{g}]$ denotes the (properly renormalized) on-shell action of the bulk field theory as a functional of prescribed boundary data $\hat{\phi}$ and $\hat{g}_{ij}$. On the left hand side, we have the generating functional for the boundary quantum field theory, which lives on a background with metric $\hat{g}_{ij}$, and whose action functional is equal to the action functional at an ultraviolet fixed point perturbed by the insertion of the relevant operator $\mathcal{O}$ with coupling $\hat{\phi}$. The expectation value on the left hand side of eqn. (1) is taken with respect

to the unperturbed fixed point action, indicated by the index 0. From eqn. (1) follow straightforwardly the correlation functions of the boundary field theory,

$$\langle \mathcal{O}(x) \rangle_{\hat{\phi}} = \frac{1}{\sqrt{\hat{g}(x)}} \frac{\delta S}{\delta \hat{\phi}(x)} , \quad (2)$$

$$\langle \mathcal{O}(x) \mathcal{O}(y) \rangle_{\hat{\phi}} = \frac{1}{\sqrt{\hat{g}(x)} \sqrt{\hat{g}(y)}} \left[-\frac{\delta^2 S}{\delta \hat{\phi}(x) \delta \hat{\phi}(y)} + \frac{\delta S}{\delta \hat{\phi}(x)} \frac{\delta S}{\delta \hat{\phi}(y)} \right] , \quad \text{etc.} \quad (3)$$

In eqns. (2) and (3), the expectation values are calculated in the perturbed boundary field theory, indicated by the index $\hat{\phi}$.

In the remainder of this section, let us give a brief outlook of the rest of the paper. We shall explain the general method in Sec. 2, which is subdivided into two parts. In subsection 2.1 we present in detail the equations of motion and the on-shell action of the bulk field theory, with special emphasis on our new gauge choice. The resulting equations of motion for the active scalars and the bulk graviton are coupled, so that, in the case of one scalar, they give rise to a third order ordinary differential equation (ODE). However, one solution is found, which guarantees the reduction of the problem to a second order ODE. The reader is referred to the appendix for matters of notation and for some relevant formulae, which are used in subsection 2.1. In subsection 2.2 we explain how the on-shell action is expressed in terms of the sources of the boundary field theory, which is essential for obtaining the correlation functions. Three cases are distinguished: operator flows involving active scalar operators of dimension $\Delta > d/2$, vev flows for these operators, and vev flows for operators of dimension $\Delta = d/2$. In the latter two, the appearance of the condensate is demonstrated. In Sec. 3 we shall apply our method to the GPPZ and Coulomb branch flows. The algebra involved, especially the factorization of the third order differential equations, is quite complex. Therefore, we had to make extensive use of symbolic algebra software (Maple). The reader is guided through the essential steps of the calculation with only the results of them given. Finally, Sec. 4 contains conclusions.

2 General Method

2.1 Equations of Motion and on-Shell Action

We consider a generic bulk field theory containing scalars coupled to gravity with the following action,

$$S = S_{\text{bulk}} + S_{\text{surf}} , \quad (4)$$

where the bulk term is given by²

$$S_{\text{bulk}} = \int d^{d+1}x \sqrt{\tilde{g}} \left[-\tilde{R} + \frac{1}{2} \tilde{g}^{\mu\nu} \partial_\mu \phi^I \partial_\nu \phi^I - V(\phi) \right] . \quad (5)$$

We assume that the potential $V(\phi)$ can be written in terms of a “super potential” $U(\phi)$ as

$$V = \frac{d}{4(d-1)} U^2 - \frac{1}{2} \left(\frac{\partial U}{\partial \phi^I} \right)^2 . \quad (6)$$

This relation holds for actions S stemming from super gravity theories, but might in fact be more general (see [32] and references therein). The surface term of the action, S_{surf} , consists of the Gibbons–Hawking term containing the second fundamental form, H , and a series of counter terms [33–36], of which we shall write explicitly only the first one, since the others do not contribute to the non-local correlation functions,

$$S_{\text{surf}} = \int d^d x \sqrt{g} [2H - U(\phi) + \dots] . \quad (7)$$

Our specific choice for the counter term is inspired by the Hamilton-Jacobi method in holography [37].

We are interested in the on-shell action for fluctuations around a given background solution. Let us write

$$\phi^I \rightarrow \phi^I + \varphi^I , \quad \tilde{g}_{\mu\nu} \rightarrow \tilde{g}_{\mu\nu} + h_{\mu\nu} ,$$

where φ^I and $h_{\mu\nu}$ are fluctuations around a solution given by ϕ^I and $\tilde{g}_{\mu\nu}$. Then, the first order variation of S_{bulk} is obtained as (N^μ is the surface normal vector)

$$\delta S_{\text{bulk}}[\phi, \tilde{g}; \varphi, h] = \int d^d x \sqrt{g} \left[N_\mu \tilde{\nabla}_\nu h_{\lambda\rho} \left(\tilde{g}^{\mu\nu} \tilde{g}^{\lambda\rho} - \tilde{g}^{\mu\lambda} \tilde{g}^{\nu\rho} \right) + N^\mu \varphi^I \partial_\mu \phi^I \right] , \quad (8)$$

where we have used the equations of motion,

$$\tilde{\nabla}^2 \phi^I + \frac{\partial V}{\partial \phi^I} = 0 , \quad (9)$$

$$\tilde{R}_{\mu\nu} - \frac{1}{2} \partial_\mu \phi^I \partial_\nu \phi^I - \frac{1}{2} \tilde{g}_{\mu\nu} \left[\tilde{R} - \frac{1}{2} \tilde{g}^{\lambda\rho} \partial_\lambda \phi^I \partial_\rho \phi^I + V(\phi) \right] = 0 , \quad (10)$$

in order to eliminate the bulk integral.

² Our conventions for the curvature tensor are $R^\mu{}_{\nu\rho\lambda} = \partial_\rho \Gamma^\mu{}_{\nu\lambda} + \Gamma^\mu{}_{\rho\sigma} \Gamma^\sigma{}_{\nu\lambda} - (\rho \leftrightarrow \lambda)$, $R_{\mu\nu} = R^\rho{}_{\mu\rho\nu}$. We have adorned with a tilde quantities belonging to $(d+1)$ space to distinguish them from those of the d -dimensional hyper surfaces.

To eqn. (8), we add the first order variation of the surface term, S_{surf} , and find

$$\delta S[\phi, \tilde{g}; \varphi, h] = \int d^d x \sqrt{g} \left[\left(\frac{\partial_r \phi^I - n^i \partial_i \phi^I}{n} - \frac{\partial U}{\partial \phi^I} \right) \varphi^I - \left(H^{ij} - H g^{ij} + \frac{1}{2} U g^{ij} \right) h_{ij} \right]. \quad (11)$$

In order to obtain eqn. (11), we have made use of the geometric relations governing hyper surfaces, which are listed in appendix A.

Eqn. (11) can be used to calculate the on-shell action to second order in the fluctuations around a given background solution, $\bar{\phi}, \bar{g}$, using the simple formula

$$S[\bar{\phi} + \varphi, \bar{g} + h] = S[\bar{\phi}, \bar{g}] + \delta S[\bar{\phi} + \varphi/2, \bar{g} + h/2; \varphi, h] + \mathcal{O}((\varphi, h)^3). \quad (12)$$

Specifically, we are interested in d -Poincaré invariant background solutions of the form

$$\bar{\phi}^I = \bar{\phi}^I(r), \quad ds^2 = dr^2 + e^{2A(r)} \eta_{ij} dx^i dx^j. \quad (13)$$

For a potential V of the form (6), such an ansatz satisfies the equations of motion (9) and (10), if [10,32,17]

$$\partial_r \bar{\phi}^I = \frac{\partial U}{\partial \phi^I}, \quad \partial_r A(r) = -\frac{1}{2(d-1)} U[\bar{\phi}(r)]. \quad (14)$$

Let us continue with linearizing the equations of motion, eqns. (9) and (10), around the background (13). First, eqn. (9) yields

$$\left(\partial_r^2 + dA' \partial_r + e^{-2A} \square \right) \varphi^I + \frac{\partial^2 V}{\partial \phi^I \partial \phi^J} \varphi^J + \frac{\partial V}{\partial \phi^I} h_{rr} - \frac{1}{2} \frac{\partial U}{\partial \phi^I} \left(\partial_r h_{rr} - \partial_r h_i^i + 2 \partial_i h_r^i \right) = 0. \quad (15)$$

Regarding our notation, $\square = \partial^i \partial_i$ and $\partial^i = \eta^{ij} \partial_j$, but $h_r^i = \bar{g}^{ij} h_{jr}$ and $h_j^i = \bar{g}^{ik} h_{kj}$.

Second, multiplying eqn. (10) by $N^\mu N^\nu$ and using the equation of Gauss, eqn. (A.6), one finds

$$R + H_j^i H_i^j - H^2 + N^\mu N^\nu \partial_\mu \phi^I \partial_\nu \phi^I - \frac{1}{2} \tilde{g}^{\mu\nu} \partial_\mu \phi^I \partial_\nu \phi^I + V(\phi) = 0, \quad (16)$$

whose linear form is

$$e^{-2A} P_j^i h_i^j + \frac{1}{2} U(\bar{\phi}) (\partial_r h_i^i - 2 \partial_i h_r^i) + V(\bar{\phi}) h_{rr} + \frac{\partial U}{\partial \phi^I} \partial_r \varphi^I + \frac{\partial V}{\partial \phi^I} \varphi^I = 0. \quad (17)$$

The symbol P_j^i denotes the transversal projector,

$$P_j^i = \partial^i \partial_j - \delta_j^i \square . \quad (18)$$

Third, the mixed components of Einstein's equation are obtained by multiplying eqn. (10) by $N^\mu \delta_i^\nu$ and using the equation of Codazzi, eqn. (A.7), with the result

$$\partial_i H - \nabla_j H_i^j - \frac{1}{2} N^\mu \partial_\mu \phi^I \partial_i \phi^I = 0 . \quad (19)$$

Linearizing eqn. (19) yields

$$\partial_r (\partial^j h_j^i - \partial^i h_j^j) + P_j^i h_r^j - \frac{1}{2} U(\bar{\phi}) \partial^i h_{rr} - \frac{\partial U}{\partial \bar{\phi}^I} \partial^i \varphi^I = 0 . \quad (20)$$

Last, the only part of the tangential components of eqn. (10) that is independent of eqns. (15), (17) and (20) is the following equation for the traceless transversal part of h_j^i ,

$$\left(\partial_r^2 + dA' \partial_r + e^{-2A} \square \right) \check{h}_j^i = 0 , \quad (21)$$

where

$$\check{h}_j^i = \frac{1}{\square^2} \Pi_{jl}^{ik} h_k^l , \quad \Pi_{jl}^{ik} = \frac{1}{2} (P^{ik} P_{jl} + P_l^i P_j^k) - \frac{1}{d-1} P_j^i P_l^k . \quad (22)$$

Eqn. (21) is identical to the equation for a free massless scalar, as is well known.

After differentiating eqn. (17) with respect to r , one can eliminate $P_j^i h_i^j$ using eqns. (17) and (20), and then use eqn. (15) to obtain

$$\begin{aligned} \frac{d}{2} U(\bar{\phi}) \partial_r h_{rr} - 2V(\bar{\phi}) h_{rr} + (d-1) e^{-2A} \square h_{rr} + 2(d-1) \frac{\partial U}{\partial \bar{\phi}^I} \partial_r \varphi^I - 2 \frac{\partial V}{\partial \bar{\phi}^I} \varphi^I \\ + (d-1) \partial_r (\partial_r h_i^i - 2\partial_i h_r^i) - U(\bar{\phi}) (\partial_r h_i^i - 2\partial_i h_r^i) = 0 . \end{aligned} \quad (23)$$

The two equations (15) and (23) depend on the three variables φ^I , h_{rr} , and $(\partial_r h_i^i - 2\partial_i h_r^i)$, the last of which can be eliminated by fixing a suitable gauge. We shall do this now.

Under a diffeomorphism, $\delta \tilde{g}_{\mu\nu} = \bar{\nabla}_\mu \xi_\nu + \bar{\nabla}_\nu \xi_\mu$, the components of the metric fluctuations transform as

$$\delta h_{ij} = \partial_i \xi_j + \partial_j \xi_i + 2A' \bar{g}_{ij} \xi_r , \quad (24)$$

$$\delta h_{ir} = \partial_i \xi_r + \partial_r \xi_i - 2A' \xi_i , \quad (25)$$

$$\delta h_{rr} = 2\partial_r \xi_r . \quad (26)$$

Thus, the combination $\partial_r h_i^i - 2\partial_i h_r^i$, which appears in the equations of motion (15) and (17), transforms as

$$\delta(\partial_r h_i^i - 2\partial_i h_r^i) = 2d \partial_r (A' \xi_r) - 2e^{-2A} \square \xi_r . \quad (27)$$

Hence, we can use first ξ_r to fix $\partial_r h_i^i - 2\partial_i h_r^i = 0$, and then ξ_i to obtain $h_r^i = 0$. Thus, our gauge is

$$\partial_r h_i^i = h_r^i = 0 . \quad (28)$$

The gauge (28) is accessible via non-singular gauge transformations and does not involve the field sources.³

In the gauge (28), the equations of motion (15), (17) and (20) become

$$\left(\partial_r^2 + dA' \partial_r + e^{-2A} \square \right) \varphi^I + \frac{\partial^2 V}{\partial \bar{\phi}^I \partial \bar{\phi}^J} \varphi^J + \frac{\partial V}{\partial \bar{\phi}^I} h_{rr} - \frac{1}{2} \frac{\partial U}{\partial \bar{\phi}^I} \partial_r h_{rr} = 0 , \quad (29)$$

$$e^{-2A} P_j^i h_i^j + V(\bar{\phi}) h_{rr} + \frac{\partial U}{\partial \bar{\phi}^I} \partial_r \varphi^I + \frac{\partial V}{\partial \bar{\phi}^I} \varphi^I = 0 , \quad (30)$$

and

$$\partial_r \partial^j h_j^i - \frac{1}{2} U(\bar{\phi}) \partial^i h_{rr} - \frac{\partial U}{\partial \bar{\phi}^I} \partial^i \varphi^I = 0 , \quad (31)$$

respectively, whereas eqn. (23) becomes

$$\frac{d}{2} U(\bar{\phi}) \partial_r h_{rr} - 2V(\bar{\phi}) h_{rr} + (d-1) e^{-2A} \square h_{rr} + 2(d-1) \frac{\partial U}{\partial \bar{\phi}^I} \partial_r \varphi^I - 2 \frac{\partial V}{\partial \bar{\phi}^I} \varphi^I = 0 . \quad (32)$$

Eqns. (29) and (32) form a system of coupled ODEs for h_{rr} and φ^I . One solution can be easily found by trial and error and is given by

$$\varphi^I = \chi \frac{\partial U}{\partial \bar{\phi}^I} , \quad h_{rr} = 2\partial_r \chi , \quad (33)$$

where χ satisfies the first order ODE

$$U(\bar{\phi}) \partial_r \chi + \partial_r U[\bar{\phi}(r)] \chi + \frac{2(d-1)}{d} e^{-2A} \square \chi = 0 . \quad (34)$$

³ In contrast, the gauge $\partial_r h_i^i - 2\partial_i h_r^i = dA' h_{rr}$, which would eliminate the terms with $\partial_r h_{rr}$ from eqn. (23), needs a singular ξ_r in the asymptotic region ($r \rightarrow \infty$). The fields would thus acquire a spurious asymptotic behaviour, and we must reject this gauge. One could also use ξ_r to eliminate h_{rr} , as was done in [28]. However, the resulting fluctuation equations appeared to be difficult to analyze.

Hence, in the case of only one active scalar, φ , the ODE (34) directly translates into a first order ODE of the form $[\partial_r + c(r)]\varphi = 0$ for the special solution given by eqn. (33), with a unique function $c(r)$. This means that the third order equation of motion for the general φ stemming from the system of equations (29) and (32) factorizes into the form

$$[\partial_r^2 + a(r)\partial_r + b(r)][\partial_r + c(r)]\varphi = 0 . \quad (35)$$

Finding the functions $a(r)$ and $b(r)$ is straightforward and is similar to the division of polynomials. This fact makes our approach feasible.

We shall now turn our attention to the on-shell action. It is straightforward to realize that, for a background of the form (13), both the background value and the first order variation of S vanish,

$$S[\bar{\phi}, \bar{g}] = 0 , \quad (36)$$

$$\delta S[\bar{\phi}, \bar{g}; \varphi, h] = 0 . \quad (37)$$

Eqn. (36) means that the normalization of the expectation values in eqns. (2) and (3) is the same as for the unperturbed theory. Moreover, it is necessary in supersymmetric settings. Eqn. (37) seems to imply that holographic one-point functions vanish. This observation has already been made by Arutyunov, Frolov and Theisen [29]. However, this fact survives only, if no additional counter terms are needed, which is the case for operator flows. We shall discuss this point in subsection 2.2.

In order to obtain the term of S quadratic in the fluctuations, we expand the second term of the right hand side of eqn. (12) and obtain, after substituting the background (13),

$$\begin{aligned} S[\bar{\phi} + \varphi, \bar{g} + h] = \int d^d x \sqrt{\bar{g}} \left[\frac{1}{2} \varphi^I \partial_r \varphi^I - \frac{1}{2} \frac{\partial^2 U}{\partial \bar{\phi}^I \partial \bar{\phi}^J} \varphi^I \varphi^J + \frac{1}{4} h_j^i \partial_r h_i^j - \frac{1}{4} h_i^i \partial_r h_j^j \right. \\ \left. - \frac{1}{4} \frac{\partial U}{\partial \bar{\phi}^I} \varphi^I (h_{rr} + h_i^i) - \frac{1}{8} U(\bar{\phi}) h_i^i h_{rr} - \frac{1}{2} h_j^i \partial_i h_r^j + \frac{1}{2} h_i^i \partial_j h_r^j \right] + \dots . \end{aligned} \quad (38)$$

It takes only a few steps to show that, by virtue of eqn. (20) and the gauge (28), eqn. (38) can be rewritten as

$$\begin{aligned} S = \frac{1}{4} \int d^d x \sqrt{\bar{g}} \left[2 \varphi^I \partial_r \varphi^I - 2 \frac{\partial^2 U}{\partial \bar{\phi}^I \partial \bar{\phi}^J} \varphi^I \varphi^J - \frac{\partial U}{\partial \bar{\phi}^I} \varphi^I h_{rr} \right. \\ \left. + h_j^i \frac{1}{\square^2} \partial_r (\Pi_{il}^{jk} h_k^l) + \frac{d}{d-1} h_j^i P_i^j \frac{1}{\square^2} \partial_r (P_l^k h_k^l) \right] . \end{aligned} \quad (39)$$

The last term on the first line will not contribute to the non-local two-point functions.

2.2 Sources and Responses

We shall begin this section introducing the reader to what we exactly mean by the terms “sources” and “responses.” In the case of scalar fields in a pure AdS background, these notions are identical to the regular and irregular boundary data, respectively [38]. In the AdS/CFT correspondence, they have been used in [39,40]. For simplicity, let us consider a single scalar field that couples to an operator of scaling dimension Δ . There are three different cases, operator flows involving scalar operators of dimension $\Delta > d/2$, vev flows involving these scalars, and vev flows involving operators of dimension $\Delta = d/2$. We shall start by analyzing in detail the operator flows, and consider the differences in the other two cases later.

The behaviour of the fields φ and h_j^i for large r can be described as [5,33,36]

$$\varphi(x, r) = e^{(\lambda-d)r/l} [\hat{\varphi}(x) + \dots] + e^{-\lambda r/l} [\check{\varphi}(x) + \dots] , \quad (40)$$

$$h_j^i(x, r) = [\hat{h}_j^i(x) + \dots] + e^{-dr/l} [\check{h}_j^i(x) + \dots] . \quad (41)$$

Here, $\lambda > d/2$, so that the first pairs of brackets on the right hand sides of eqns. (40) and (41) denote the leading series. The coefficients $\hat{\varphi}(x)$ and $\hat{h}_j^i(x)$ are regarded as the sources, and the remaining terms in the brackets depend locally on them.⁴ The second pairs of brackets denote the subleading series, which are determined by the responses $\check{\varphi}(x)$ and $\check{h}_j^i(x)$.⁵ The general solution to the coupled equations of motion, eqns. (29) and (32), is determined uniquely by specifying three integration constants, say, $\hat{\varphi}$, $P_j^i \hat{h}_i^j$ and $\check{\varphi}$, but it is in general not regular in the bulk. Thus, imposing one suitable regularity condition, only two constants need to be specified, which we choose to be the sources, $\hat{\varphi}$ and $P_j^i \hat{h}_i^j$. Then, the responses, $\check{\varphi}$ and $P_j^i \check{h}_i^j$, are uniquely determined and give rise to the holographic two-point functions. It seems clear that, since only one condition can be imposed, one cannot demand regularity for both fields, φ and h_j^i . Notice that, in general, each of the responses will depend on both sources. This is not so in the pure AdS background, where the scalar and gravity equations of motion decouple.

⁴ There is another way of defining the sources, namely by imposing Dirichlet boundary conditions on the fields at some cut-off boundary. The boundary values are the bare sources, which have to be rescaled when sending the cut-off to infinity. This method has been used successfully in the AdS/CFT correspondence, but it is technically very difficult to solve a Dirichlet boundary value problem in the RG flow backgrounds considered here.

⁵ Generically, the source and response series are in powers of $e^{-2r/l}$ with coefficients that depend locally on the sources and responses, respectively. However, for $\lambda = d/2 + k$, with positive integer k , as well as for even d , the source series contain terms of the form $r/l e^{-2nr/l}$ with $n \geq k$ and $n \geq d/2$, respectively.

In order to analyze the equations of motion in the asymptotically AdS region of the bulk (for large r), we let $A(r) = r/l + \dots$. Moreover, we need the following expressions of the potentials U and V close to the UV fixed point [41],

$$U(\phi) = -\frac{2(d-1)}{l} + \frac{\lambda-d}{2l}\phi^2 + \dots, \quad (42)$$

$$V(\phi) = \frac{d(d-1)}{l^2} - \frac{1}{2}m^2\phi^2 + \dots, \quad (43)$$

where $m^2 l^2 = \lambda(\lambda-d)$. Since the background field satisfies eqn. (14), eqn. (42) tells us that $\bar{\phi}$ behaves as

$$\bar{\phi}(r) = \hat{\phi} e^{(\lambda-d)r/l} + \dots. \quad (44)$$

Using eqn. (30) one can determine the asymptotic behaviour of h_{rr} ,

$$h_{rr}(x, r) = -\frac{l^2 e^{-2r/l}}{d(d-1)} P_j^i \hat{h}_i^j + \frac{\lambda-d}{d-1} \hat{\phi} \hat{\phi} e^{2(\lambda-d)r/l} + \dots + \frac{2m^2 l^2}{d(d-1)} \hat{\phi} \check{\phi} e^{-dr/l} + \dots. \quad (45)$$

Then, from eqn. (31) one finds the asymptotic behaviour of $P_j^i h_i^j$,

$$P_j^i h_i^j(x, r) = P_j^i \hat{h}_i^j(x) - \frac{l^2}{2d} e^{-2r/l} \square P_j^i \hat{h}_i^j + \dots + \frac{(\lambda-d)(2\lambda-d)}{d^2} \hat{\phi} \square \check{\phi} e^{-dr/l} + \dots. \quad (46)$$

It is reassuring that the term with $e^{2(\lambda-d)r/l}$ vanishes, and that one reproduces the behaviour of eqn. (41).

The traceless transversal part of h_j^i obeys the second order ODE (21). The asymptotic behaviour of the solution is given by

$$\check{h}_j^i(x, r) = \hat{h}_j^i(x) + \frac{l^2}{2(d-2)} e^{-2r/l} \square \hat{h}_j^i(x) + \dots + \check{h}_k^l(x) e^{-dr/l} + \dots. \quad (47)$$

As in the pure AdS background, the response $\check{h}_k^l$ is determined in terms of the source $\hat{h}_j^i$ by imposing the condition that the solution be regular in the bulk.

Let us now return to the on-shell action, eqn. (39). After inserting the expansions (40), (46) and (47), the finite terms give us the generating functional for two-point functions of the boundary field theory. However, we must first look at the divergent terms. The leading divergent terms (proportional to $e^{dr/l}$ and $e^{(2\lambda-d)r/l}$) should vanish, since the appropriate counter terms have already been added. In eqn. (39), there are no terms proportional to $e^{dr/l}$, while it is

easy to see that the terms proportional to $e^{(2\lambda-d)r/l}\hat{\varphi}^2$ cancel. Moreover, the remaining divergent terms must be local, so that they can be cancelled by adding further local counter terms (involving the curvature and derivatives of ϕ). The two terms on the second line of eqn. (39) might worry us, since they are non-local in the sources. However, from eqns. (46) and (47) follows

$$\begin{aligned} h_j^i \frac{1}{\square^2} \partial_r \left(\Pi_{il}^{jk} h_k^l \right) + \frac{d}{d-1} h_j^i P_i^j \frac{1}{\square^2} \partial_r (P_l^k h_k^l) \\ = -\frac{l}{d-2} \hat{h}_j^i \frac{\Pi_{il}^{jk}}{\square} \hat{h}_k^l e^{-2r/l} + \frac{l}{d-1} \hat{h}_j^i \frac{P_i^j P_l^k}{\square} \hat{h}_k^l e^{-2r/l} + \dots, \end{aligned} \quad (48)$$

and the non-local terms (stemming from the terms of the projectors with four distinct indices) cancel. Hence, the remaining divergent terms can be cancelled by adding appropriate local counter terms. These counter terms will contribute further contact terms to the correlators, but we will not consider them explicitly.

We can now turn our attention to the finite, non-local part of the action. Inserting eqns. (40), (46) and (47) into eqn. (39), we obtain

$$S_{\text{fin}} = \frac{1}{l} \int d^d x \left[-\left(\lambda - \frac{d}{2} \right) \hat{\varphi} \check{\varphi} - \frac{d}{4} \hat{h}_j^i \frac{\Pi_{il}^{jk}}{\square^2} \check{h}_k^l - \frac{d^2}{4(d-1)} \hat{h}_j^i \frac{P_i^j P_l^k}{\square^2} \check{h}_k^l \right]. \quad (49)$$

From S_{fin} , correlation functions are obtained using the correspondence formula (1), as in eqns. (2) and (3). When doing so, we shall multiply S_{fin} by a constant of dimension length, κ , in order to make the exponent dimensionless.

The first term in eqn. (49) is responsible for the scalar two-point function, $\langle \mathcal{O} \mathcal{O} \rangle$. From the AdS/CFT correspondence it is known that the boundary field theory operators have dimension $\Delta = \lambda > d/2$ at the conformal fixed point. The second and third terms give the two-point function of the energy momentum tensor, $\langle T_j^i T_l^k \rangle$. However, since the scalar φ couples to $P_j^i h_i^j$, the response $\check{\varphi}$ will also depend on the source $\hat{h}_j^i$, and, *vice versa*, the response $P_j^i \check{h}_i^j$ will depend on the source $\hat{\varphi}$. Thus, the first and third terms of eqn. (49) give also rise to a two-point function of the form $\langle T_j^i \mathcal{O} \rangle$. This two-point function satisfies the Ward identity for $4d$ Poincaré invariance,

$$\frac{\partial}{\partial x^i} \langle T_j^i(x) \mathcal{O}(y) \rangle = 0, \quad (50)$$

since it is proportional to the transversal projector P_j^i .

Let us consider now the case $\lambda < d/2$, which corresponds to vev flows. The formulae (40) to (47) remain valid as before, but the interpretation of the fields $\hat{\varphi}$ and $\check{\varphi}$ changes. In fact, since $\lambda - d < -\lambda$, the series with $\check{\varphi}$ in eqn. (40) becomes the leading one. Correspondingly, we interpret $\check{\varphi}$ as the source,

whereas $\hat{\varphi}$ is the response. The background, behaving as in eqn. (44), follows the subleading behaviour of the fluctuations. This means that the background source is zero, *i.e.*, there is no operator insertion in the boundary field theory action.

The on-shell action, eqn. (39), becomes

$$S = -\frac{1}{l} \left(\lambda - \frac{d}{2} \right) \int d^d x \left(\hat{\varphi} \check{\varphi} + \check{\varphi}^2 e^{(d-2\lambda)r/l} + \dots \right) , \quad (51)$$

where we have not written those terms that remain unchanged with respect to eqn. (49). However, since we now have $d > 2\lambda$, the second term is divergent (in this case, the counter term with $U(\phi)$ cancels the sub-leading term), and we need to add a counter term,

$$S_{\text{ct}} = \frac{1}{l} \left(\lambda - \frac{d}{2} \right) \int d^d x \sqrt{g} \phi^2 . \quad (52)$$

Notice that the counter term must be expressed in terms of the full fields, because of the universality of the action. Adding eqn. (51) to (52), we find the finite on-shell action,

$$S_{\text{fin}} = \frac{1}{l} \left(\lambda - \frac{d}{2} \right) \int d^d x \left(\check{\varphi} \hat{\varphi} + 2\check{\varphi} \hat{\hat{\phi}} + h_i^i \hat{\hat{\phi}} \check{\varphi} + \dots \right) . \quad (53)$$

Eqn. (53) gives correlation functions of operators, whose dimension at the conformal fixed point is $\Delta = d - \lambda > d/2$. In fact, for the AdS background, eqns. (49) and (53) are identical, when λ is expressed in terms of Δ . For RG flow backgrounds, we explicitly obtain the condensate

$$\langle \mathcal{O}(x) \rangle = (d - 2\Delta) \frac{\kappa}{l} \hat{\hat{\phi}} . \quad (54)$$

The contact contribution to $\langle T_j^i \mathcal{O} \rangle$ from the third term on the right hand side of eqn. (53) has its origin in the fact that T_j^i is the energy-momentum operator of the boundary field theory including the source term [42].

Let us finally consider the special case of operators with scaling dimension $\lambda = d/2$, which also corresponds to vev flows. While the background field behaves asymptotically as

$$\bar{\phi}(r) = e^{-dr/(2l)} \check{\phi} + \dots , \quad (55)$$

the leading behaviour of the perturbation φ is

$$\varphi(r, x) = e^{-dr/(2l)} \left[\frac{r}{l} \hat{\varphi}(x) + \check{\varphi}(x) \right] + \dots . \quad (56)$$

Again, the background behaves as the subleading part of the fluctuation. Hence, the background source is zero, and the RG flow must be a vev flow.

Inserting eqn. (56) into the action (39), one finds

$$S = \frac{1}{2l} \int d^d x \left(\frac{r}{l} \hat{\varphi}^2 + \hat{\varphi} \check{\varphi} + \dots \right) , \quad (57)$$

where we have not written those terms that remain unchanged with respect to eqn. (49). The term with r/l in S is logarithmically divergent and must be cancelled by adding a suitable counter term,⁶

$$S_{\text{ct}} = -\frac{1}{2r} \int d^d x \sqrt{g} \phi^2 . \quad (58)$$

As before, the counter term must be expressed in terms of the full fields ϕ and g_{ij} , not only in terms of the fluctuations φ and h_j^i , since it should be valid for any background solution. Adding S_{ct} to eqn. (57) yields the finite action

$$S_{\text{fin}} = -\frac{1}{l} \int d^d x \left(\frac{1}{2} \hat{\varphi} \check{\varphi} + \hat{\varphi} \check{\check{\varphi}} + \frac{1}{2} \hat{\varphi} \check{\check{\varphi}} \hat{h}_i^i + \dots \right) , \quad (59)$$

where the ellipses indicate the last two terms of eqn. (49), which remain unchanged. One can check easily that the formula (59) yields the correct (*i.e.*, positive definite) conformal two-point function $\langle \mathcal{O}(x) \mathcal{O}(y) \rangle$ in the case of the AdS background. The scalar operators have dimension $\Delta = d/2$ at the conformal fixed point.

The second term in eqn. (59) yields the condensate of the boundary field theory,

$$\langle \mathcal{O}(x) \rangle = -\frac{\kappa}{l} \check{\check{\varphi}} , \quad (60)$$

whereas the third term again gives a local contribution to the correlator $\langle T_j^i \mathcal{O} \rangle$.

3 Particular Flow Solutions

3.1 GPPZ Flow

An explicit flow solution describing the renormalization group flow from $\mathcal{N} = 4$ SYM theory in the UV to $\mathcal{N} = 1$ SYM theory in the IR ($d = 4$) was

⁶ “Logarithmically” because r/l is proportional to the logarithm of the volume of the boundary space.

found by GPPZ [7]. The flow is generated by perturbing the SYM theory action by inserting an operator of dimension $\Delta = 3$. The super potential $U(\phi)$ responsible for the GPPZ flow is, in our notation (only the non-zero active scalar is shown),

$$U(\phi) = -\frac{3}{l} \left(\cosh \frac{\phi}{\sqrt{3}} + 1 \right) , \quad (61)$$

and the background flow solution is given by

$$\bar{\phi}(r) = \sqrt{3} \ln \frac{1 + e^{-r/l}}{1 - e^{-r/l}} , \quad (62)$$

$$A(r) = \frac{1}{2} \left[\frac{r}{l} + \ln \left(e^{r/l} - e^{-r/l} \right) \right] . \quad (63)$$

Towards the horizon, the background field $\bar{\phi}$ behaves as

$$\bar{\phi}(r) = 2\sqrt{3} e^{-r/l} + \dots , \quad (64)$$

so that, according to eqn. (44), $\Delta = \lambda = 3$ and $\hat{\phi}(x) = 2\sqrt{3}$.

In order to study the fluctuation equations, it is advantageous to introduce the variable

$$v = 1 - e^{-2r/l} . \quad (65)$$

The system of equations (29) and (32) is equivalent to a third order ODE for φ , which is of the form [cf. eqn. (35)]

$$\left[\frac{d^2}{dv^2} + a(v) \frac{d}{dv} + b(v) \right] \left[\frac{d}{dv} - \frac{1}{2(1-v)} + \frac{p^2 l^2}{8} \right] \varphi(v) = 0 . \quad (66)$$

The first order factor is obtained from eqn. (34), and the coefficients $a(v)$ and $b(v)$ have been found using symbolic algebra software (Maple),

$$a(v) = \frac{p^2 l^2 v^2 + 8v - 2p^2 l^2 v + 36}{v(p^2 l^2 v^2 + 4v - p^2 l^2 v + 12)} , \quad (67)$$

$$b(v) = \frac{p^4 l^4 v(1-v)^2 + p^2 l^2 (1-v)(v^2 + 4v - 16) + 4(v^2 + 5v - 2)}{4v(p^2 l^2 v^2 + 4v - p^2 l^2 v + 12)(1-v)^2} . \quad (68)$$

The solution of the second order ODE,

$$\left[\frac{d^2}{dv^2} + a(v) \frac{d}{dv} + b(v) \right] \psi(v) = 0 , \quad (69)$$

which is regular at $v = 0$, is

$$\psi(v) = C_2 \sqrt{1-v} \left\{ 4(1+v) F\left(\frac{3+\alpha}{2}, \frac{3-\alpha}{2}; 3; v\right) + 8 F\left(\frac{1+\alpha}{2}, \frac{1-\alpha}{2}; 2; v\right) \right\} , \quad (70)$$

where $\alpha = \sqrt{1-p^2 l^2}$, and C_2 is a constant.

Now, one has to solve the inhomogeneous first order equation

$$\left[\frac{d}{dv} + \frac{5 - \alpha^2 - v(1 - \alpha^2)}{8(1-v)} \right] \varphi(v) = \psi(v) . \quad (71)$$

Let us write $\varphi(v)$ as

$$\varphi(v) = \sqrt{1-v} \left[C_1 e^{-(1-\alpha^2)v/8} + \frac{64C_2}{1-\alpha^2} F\left(\frac{1+\alpha}{2}, \frac{1-\alpha}{2}; 2; v\right) + C_2 \lambda(v) \right] , \quad (72)$$

where the term with the constant C_1 is the universal solution of eqn. (71). Now, we are left with the simpler equation for $\lambda(v)$,

$$\left[\frac{d}{dv} - \frac{1 - \alpha^2}{8} \right] \lambda(v) = -4(1-v) F\left(\frac{3+\alpha}{3}, \frac{3-\alpha}{2}; 3; v\right) , \quad (73)$$

but we will not have to solve it explicitly.

Instead, we shall look at the asymptotic behaviour of the fields at the horizon. First, let us introduce

$$u = \sqrt{1-v} , \quad (74)$$

so that the horizon is at $u = 0$. Then, we can describe the asymptotic behaviour of the field φ by the expansion

$$\varphi(u) = u \left[A_1 + \ln u (A_2 u^2 + A_3 u^4 + A_4 u^6 + \dots) + B_1 u^2 + B_2 u^4 + B_3 u^6 + \dots \right] . \quad (75)$$

This expansion uniquely determines φ , when the three coefficients A_1 , A_2 and B_1 are specified. The higher order coefficients can be obtained recursively from the third order ODE (66). We have used Maple to perform this task.

The solution (72) has only two coefficients, C_1 and C_2 , since we have used only the solution of the second order ODE (69) that is regular in the bulk. After changing variables to u , one can see from eqn. (73) that $\lambda(u)$ is of order $u^4 \ln u$. Thus, the coefficients A_1 , A_2 and B_1 are related to C_1 and C_2 by (see

[43] for the asymptotic expansion of the hypergeometric function)

$$A_1 = C_1 + \frac{64\tilde{C}_2}{1 - \alpha^2} , \quad (76)$$

$$A_2 = 32\tilde{C}_2 , \quad (77)$$

$$B_1 = \frac{1 - \alpha^2}{8} C_1 + 16\tilde{C}_2 \left[\psi\left(\frac{3 + \alpha}{2}\right) + \psi\left(\frac{3 - \alpha}{2}\right) - \psi(1) - \psi(2) \right] . \quad (78)$$

Here, we have introduced for brevity of notation

$$\tilde{C}_2 = C_2 \left[\Gamma\left(\frac{3 + \alpha}{2}\right) \Gamma\left(\frac{3 - \alpha}{2}\right) \right]^{-1} .$$

Obviously, A_1 is identified with the source, $\hat{\varphi}$, whereas B_1 is the response, $\check{\varphi}$. Specifying the source $P_j^i \hat{h}_i^j$ determines also A_2 uniquely. In fact, $P_j^i \hat{h}_i^j$ is obtained from eqn. (30), where h_{rr} is to be expressed in terms of φ by solving algebraically eqns. (29) and (32). After the necessary algebra we find

$$P_j^i \hat{h}_i^j(u, p) = \frac{\sqrt{3}}{l^2} [A_1(\alpha^2 - 1) + 2A_2] + \mathcal{O}(u^2) . \quad (79)$$

Hence, after determining A_1 and A_2 in terms of the sources $\hat{\varphi}$ and $P_j^i \hat{h}_i^j$, solving eqns. (76) and (77) for C_1 and $\tilde{C}_2$, and substituting the solution into eqn. (78), we obtain

$$\check{\varphi}(p) = \frac{p^2 l^2}{4} f_1(p^2) \hat{\varphi}(p) + \frac{\sqrt{3} l^2}{12} \left[f_1(p^2) - \frac{1}{2} \right] P_j^i \hat{h}_i^j(p) , \quad (80)$$

where the function $f_1(p^2)$ is

$$f_1(p^2) = \psi\left(\frac{3 + \sqrt{1 - p^2 l^2}}{2}\right) + \psi\left(\frac{3 - \sqrt{1 - p^2 l^2}}{2}\right) - \psi(2) - \psi(1) . \quad (81)$$

It remains to determine the response $P_j^i \check{h}_i^j$. We proceed as for the derivation of eqn. (79), but do not truncate at the lowest order. It is necessary to substitute the recursively determined coefficients A_3 , A_4 , B_2 and B_3 , after which one finds

$$\begin{aligned} P_j^i \check{h}_i^j(u, p) &= P_j^i \hat{h}_i^j(p) + u^2 \frac{p^2 l^2}{8} P_j^i \hat{h}_i^j(p) \\ &+ u^4 \left\{ \frac{\sqrt{3} p^4 l^2}{16} \left[f_1(p^2) + \frac{1}{2} \right] \hat{\varphi}(p) + \frac{p^2 l^2}{16} \left[f_1(p^2) + \frac{p^2 l^2 + 4}{8} \right] P_j^i \hat{h}_i^j(p) \right\} + \dots . \end{aligned} \quad (82)$$

Here, we have not explicitly written the term of order $u^4 \ln u$, whose coefficient is a local function of the sources. One can easily check that eqn. (82) agrees

with the general form (46) up to terms of order u^4 that depend locally on the sources. The term proportional to u^4 is the response $P_j^i \check{h}_i^j$.

For completeness, we shall repeat here the solution for the traceless transversal part of h_j^i [27,28,30]. In terms of the variable v , eqn. (21) reads

$$\left[v(1-v) \frac{d^2}{dv^2} + (2-v) \frac{d}{dv} - \frac{p^2 l^2}{4} \right] \check{h}_j^i(v) = 0 , \quad (83)$$

and its solution, which is regular at $v = 0$, is

$$\check{h}_j^i(v) = C_j^i F \left(\frac{\sqrt{-p^2 l^2}}{2}, -\frac{\sqrt{-p^2 l^2}}{2}; 2; v \right) . \quad (84)$$

After asymptotically expanding the hypergeometric function for small $u = \sqrt{1-v}$ [43], eqn. (84) can be expressed as

$$\check{h}_j^i(u) = \hat{h}_j^i(p) \left\{ 1 - \frac{p^2 l^2}{4} u^2 - \frac{p^2 l^2 (4 + p^2 l^2)}{32} \left[2 \ln u + f_2(p^2) + \frac{1}{2} \right] u^4 + \dots \right\} , \quad (85)$$

where we have introduced the source $\hat{h}_j^i$, and $f_2(p^2)$ is

$$f_2(p^2) = \psi \left(2 - \frac{\sqrt{-p^2 l^2}}{2} \right) + \psi \left(2 + \frac{\sqrt{-p^2 l^2}}{2} \right) - 2\psi(2) . \quad (86)$$

The term in eqn. (85) proportional to u^4 is the response $\check{h}_j^i$.

We are now in a position to substitute the responses, $\check{\varphi}$, $P_j^i \check{h}_i^j$ and $\check{h}_j^i$, into the expression for the finite part of the on-shell action, eqn. (49). One obtains

$$S_{\text{fin}} = \int \frac{d^4 p}{(2\pi)^4} \left\{ -\hat{\varphi}(-p) \frac{p^2 l}{4} f_1(p^2) \hat{\varphi}(p) - \hat{\varphi}(-p) \frac{\sqrt{3}l}{6} f_1(p^2) P_j^i \hat{h}_i^j(p) \right. \\ \left. - \hat{h}_j^i(-p) \frac{l}{4p^2} \left[\frac{1}{3} \left(f_1(p^2) + \frac{4 + p^2 l^2}{8} \right) P_i^j P_k^l - \frac{4 + p^2 l^2}{8} \left(f_2(p^2) + \frac{1}{2} \right) \Pi_{ik}^{jl} \right] \hat{h}_l^k(p) \right\} . \quad (87)$$

Finally, from eqn. (87) one can read off the two-point functions

$$\langle \mathcal{O}(p) \mathcal{O}(-p) \rangle = \frac{\kappa l}{2} p^2 f_1(p^2) , \quad (88)$$

$$\langle T_j^i(p) \mathcal{O}(-p) \rangle = \frac{\sqrt{3} \kappa l}{3} f_1(p^2) P_j^i , \quad (89)$$

$$\langle T_j^i(p) T_l^k(-p) \rangle = \frac{2 \kappa l}{p^2} \left[-\frac{(4 + p^2 l^2)}{8} f_2(p^2) \Pi_{jl}^{ik} + \frac{1}{3} f_1(p^2) P_j^i P_l^k \right] . \quad (90)$$

In passing from eqn. (87) to eqn. (90) we have omitted a local term proportional to $(4 + p^2 l^2)$. The correlator (88) and the traceless transversal part of the correlator (90) agree with the expressions known from the literature up to normalization [27,29,30]. Taking the trace of the energy momentum tensors, we find the following relations,

$$\langle T_i^i(p) \mathcal{O}(-p) \rangle = \sqrt{3} \kappa l p^2 f_1(p^2) = 2\sqrt{3} \langle \mathcal{O}(p) \mathcal{O}(-p) \rangle , \quad (91)$$

$$\langle T_i^i(p) T_j^j(-p) \rangle = 6 \kappa l p^2 f_1(p^2) = 12 \langle \mathcal{O}(p) \mathcal{O}(-p) \rangle , \quad (92)$$

leading to the conclusion that

$$T_i^i = 2\sqrt{3} \mathcal{O} . \quad (93)$$

This relation is direct evidence of the expected operator relation

$$T_i^i(x) = \beta^n \mathcal{O}_n(x) , \quad (94)$$

where β^n are the beta functions corresponding to the couplings of the primary operators $\mathcal{O}_n$ [44]. For the active scalar operator in the GPPZ flow, we find

$$\beta = 2\sqrt{3} = -(\Delta - d) \hat{\phi}(x) , \quad (95)$$

which agrees with the renormalized holographic beta function [27,41].

3.2 Coulomb Branch Flow

In this section, we shall consider as an example one of the five renormalization group flows involving a single active scalar from the **20'** of $SO(6)$ in $d = 4$, $\mathcal{N} = 4$ SYM theory [11,45]. It was called $n = 2$ flow in [11]. These flows are vev flows moving the field theory onto the Coulomb branch.

In our notation, the potential $U(\phi)$ responsible for the Coulomb branch flow is given by

$$U(\phi) = -\frac{4}{l} \left(e^{-\phi/\sqrt{6}} + \frac{1}{2} e^{2\phi/\sqrt{6}} \right) . \quad (96)$$

Introducing the new variable

$$v = e^{\sqrt{6}\bar{\phi}(r)/2} , \quad (97)$$

the various quantities can be re-expressed as

$$\begin{aligned} U(\bar{\phi}) &= -\frac{2(v+2)}{lv^{1/3}}, \quad \frac{\partial U}{\partial \bar{\phi}} = \frac{4(1-v)}{\sqrt{6}lv^{1/3}}, \\ V(\bar{\phi}) &= \frac{4(1+2v)}{l^2v^{2/3}}, \quad \frac{\partial V}{\partial \bar{\phi}} = -\frac{8(1-v)}{\sqrt{6}l^2v^{2/3}}, \quad \frac{\partial^2 V}{\partial \bar{\phi}^2} = \frac{4(2+v)}{3l^2v^{2/3}}, \\ e^{2A(r)} &= \frac{l^2v^{2/3}}{L^2(1-v)}, \quad \frac{dv}{dr} = \frac{2}{l}v^{2/3}(1-v), \end{aligned} \quad (98)$$

where the length L is related to the radius of $D3$ -branes, l_{D3} , by $l_{D3} = l^2/L$ [11].

The horizon is at $v = 1$. The asymptotic behaviour of the background field $\bar{\phi}$ towards the horizon can be obtained from eqns. (97) and (98) (remember that $A(r) \rightarrow r/l$),

$$\bar{\phi}(r) = -\frac{\sqrt{6}l^2}{3L^2}e^{-2r/l} + \dots, \quad (99)$$

and we can read off $\Delta = 2$ and $\check{\bar{\phi}}(x) = -\sqrt{6}l^2/(3L^2)$.

The system of equations (29) and (32) is equivalent to a third order ODE for φ , which is of the form

$$\left[\frac{d^2}{dv^2} + a(v)\frac{d}{dv} + b(v) \right] \left[\frac{d}{dv} + \frac{1}{2+v} + \frac{1}{1-v} + \frac{3L^2p^2}{8v(2+v)} \right] \varphi(v) = 0. \quad (100)$$

Again, we have performed the factorization using Maple and obtained the coefficients $a(v)$ and $b(v)$ as

$$a(v) = \frac{5p^2L^2v^2 + 48v - 8p^2L^2 + 64}{v(2+v)[p^2L^2(v-1) + 8]}, \quad (101)$$

$$b(v) = \frac{p^4L^4(v-1)(v+2) + 4p^2L^2(4v^3 - 9v^2 + 8) + 64(v-1)(3v+2)}{4(v-1)v^2(2+v)[p^2L^2(v-1) + 8]}. \quad (102)$$

Let us now solve the second order equation,

$$\left[\frac{d^2}{dv^2} + a(v)\frac{d}{dv} + b(v) \right] \psi(v) = 0, \quad (103)$$

With the substitution

$$\psi(v) = (2+v)^{-1}v^{\alpha/2-2}\rho(v), \quad (104)$$

where the constant α is

$$\alpha = \sqrt{1 + p^2 L^2} + 1 , \quad (105)$$

eqn. (103) becomes

$$\begin{aligned} v(1-v) \left(v - \frac{(\alpha+2)(\alpha-4)}{\alpha(\alpha-2)} \right) \frac{d^2 \rho}{dv^2} \\ + \left[-(\alpha-1)v^2 + \frac{(\alpha+2)(\alpha-4)}{\alpha-2}v + (\alpha-1)v - \frac{(\alpha+2)(\alpha-4)}{\alpha-2} \right] \frac{d\rho}{dv} \\ - \frac{1}{4} [(\alpha-2)^2 v - \alpha(\alpha-4)] \rho = 0 . \end{aligned} \quad (106)$$

The solution of eqn. (106) that is regular at $v = 0$ is

$$\rho(v) = C \left[F \left(\frac{\alpha}{2} - 1, \frac{\alpha}{2} - 1; \alpha; v \right) + \frac{2}{\alpha} F \left(\frac{\alpha}{2} - 1, \frac{\alpha}{2}; \alpha; v \right) \right] , \quad (107)$$

where C is a constant.

It remains to extract the asymptotic behaviour of $\varphi(x)$ near the horizon from the first order differential equation

$$\left[\frac{d}{dv} + \frac{1}{2+v} + \frac{1}{1-v} + \frac{3L^2 p^2}{8v(2+v)} \right] \varphi(v) = \psi(v) . \quad (108)$$

Let us introduce the new variable u by $v = 1 - (l^2/L^2)u$, so that $u \approx e^{-2r/l}$ for large r . Then, the field φ has an asymptotic expansion

$$\varphi(u) = \ln u (A_1 u + A_2 u^2 + A_3 u^3 + \dots) + B_1 u + B_2 u^2 + B_3 u^3 + \dots . \quad (109)$$

Since $\varphi(u)$ satisfies the third order differential equation (100), specifying the three coefficients A_1 , B_1 and B_2 determines the field uniquely. The higher order coefficients can be determined recursively, which we have done using Maple. The coefficients A_1 and B_1 are related to the source and response, respectively, by

$$\hat{\varphi} = -2A_1, \quad \check{\varphi} = B_1 . \quad (110)$$

In order to determine the coefficient B_2 , one has to look at the leading order of the field $P_j^i h_i^j$. It is obtained from eqn. (30), where h_{rr} is to be expressed in terms of φ by solving algebraically eqns. (29) and (32). After the necessary algebra we find

$$P_j^i h_i^j(u, p) = \frac{\sqrt{6}}{l^2} \left\{ [16 - 6\alpha(\alpha-2)]A_1 + [4 + 3\alpha(\alpha-2)]B_1 + 24 \frac{L^2}{l^2} B_2 \right\} + \mathcal{O}(u) . \quad (111)$$

The additional relation between A_1 , B_1 and B_2 is obtained from eqn. (108) as follows. After changing the variable to u and inserting the asymptotic expansion (109), the left hand side of eqn. (108) reads (the recursively determined higher order coefficients must also be substituted)

$$\begin{aligned}
& -\frac{L^2}{l^2}A_1 - \frac{1}{8}\alpha(\alpha-2)A_1u \ln u \\
& + \left\{ \left[-\frac{1}{3} - \frac{1}{4}\alpha(\alpha-2) \right] A_1 + \left[\frac{1}{3} + \frac{1}{8}\alpha(\alpha-2) \right] B_1 - \frac{L^2}{l^2}B_2 \right\} u + \mathcal{O}(u^2 \ln u) ,
\end{aligned} \tag{112}$$

whereas the asymptotic expansion of the right hand side is [43]

$$\begin{aligned}
& \frac{2\Gamma(\alpha)C}{3\Gamma(\alpha/2+1)^2} \left\{ 1 + \frac{l^2}{8L^2}\alpha(\alpha-2)u \ln u \right. \\
& \left. + \frac{l^2u}{8L^2} \left[\frac{32}{8} + \alpha(\alpha-2) \left(\ln \frac{l^2}{L^2} + 2\psi\left(\frac{\alpha}{2}\right) - 2\psi(1) - 2 \right) \right] \right\} + \mathcal{O}(u^2 \ln u) .
\end{aligned} \tag{113}$$

Comparing the constant terms of eqns. (112) and (113) yields C in terms of A_1 , and the u term gives the desired relation between the three coefficients A_1 , B_1 and B_2 . Expressing A_1 , B_1 and B_2 in terms of the sources $\hat{\varphi}$ and $P_j^i \hat{h}_i^j$ and the response $\check{\varphi}$, one finds the following relation,

$$\check{\varphi}(p) = \left[-\psi\left(\frac{\alpha}{2}\right) + \psi(1) - \frac{1}{2} \ln \frac{l^2}{L^2} + \frac{4}{3\alpha(\alpha-2)} \right] \hat{\varphi}(p) + \frac{\sqrt{6}l^2}{9\alpha(\alpha-2)} P_j^i \hat{h}_i^j(p) . \tag{114}$$

It remains to determine the response $P_j^i \check{h}_i^j$. We proceed as for the derivation of eqn. (111), but do not truncate at the lowest order. It is necessary to substitute the recursively determined higher order coefficients, after which one finds

$$\begin{aligned}
P_j^i h_i^j(u, p) &= P_j^i \hat{h}_i^j(p) + u \frac{p^2 l^2}{8} P_j^i \hat{h}_i^j(p) \\
&+ u^2 \left[\frac{\sqrt{6} p^2 l^2}{24 L^2} \hat{\varphi}(p) + \frac{p^2 l^4}{384 L^2} (3 p^2 L^2 + 32) P_j^i \hat{h}_i^j(p) \right] + \dots .
\end{aligned} \tag{115}$$

Again, we have not written explicitly the term $u^2 \ln u$, whose coefficient depends locally on the sources. The u^2 term in eqn. (115) is the response $P_j^i \check{h}_i^j$.

For completeness, we shall give here also the solution for the traceless transversal part of the graviton. In terms of v , eqn. (21) reads

$$\left[v(1-v) \frac{\partial^2}{\partial v^2} + (2-v) \frac{\partial}{\partial v} - \frac{1}{4v} \alpha(\alpha-2) \right] \check{h}_j^i = 0 , \tag{116}$$

whose solution, which is regular for $v = 0$, is

$$\check{h}_j^i = C_j^i v^{\alpha/2-1} F\left(\frac{\alpha}{2} - 1, \frac{\alpha}{2} - 1; \alpha; v\right) . \quad (117)$$

Asymptotically expanding eqn. (117) for small u yields [43]

$$\check{h}_j^i(u, p) = \hat{h}_j^i(p) \left\{ 1 - \frac{p^2 l^2}{4} u - \frac{p^4 l^4}{32} u^2 \left[\ln u + \ln \frac{l^2}{L^2} + 2\psi\left(\frac{\alpha}{2}\right) - 2\psi(1) + \frac{4}{p^2 L^2} - \frac{3}{2} \right] + \dots \right\} . \quad (118)$$

The term proportional to u^2 is the response $\check{h}_j^i$. Notice that, in contrast to the massless scalar field, we cannot discard the local terms in eqn. (118), because there is a factor $1/p^4$ in the on-shell action.

Finally, we substitute the responses $\check{\varphi}$, $P_j^i \check{h}_i^j$ and $\check{h}_j^i$ into the expression for the finite on-shell action, eqns. (49) and (59). We obtain

$$\begin{aligned} S_{\text{fin}} = \int \frac{d^4 p}{(2\pi)^4} \left\{ \hat{\varphi}(-p) \frac{\sqrt{6}l}{3L^2} \delta(p) + \frac{1}{2l} \hat{\varphi}(-p) \left[\psi\left(\frac{\alpha}{2}\right) - \psi(1) + \frac{1}{2} \ln \frac{l^2}{L^2} - \frac{4}{3p^2 L^2} \right] \hat{\varphi}(p) \right. \\ - \frac{\sqrt{6}l}{3L^2} \hat{\varphi}(-p) \left(\frac{P_j^i}{3p^2} - \frac{1}{2} \delta_j^i \right) \hat{h}_i^j(p) - \hat{h}_j^i(-p) \frac{l^3}{3} P_i^j P_l^k \left(\frac{1}{32} + \frac{1}{3p^2 L^2} \right) \hat{h}_k^l(p) \\ \left. + \hat{h}_j^i(-p) \frac{l^3}{16} \Pi_{il}^{jk} \left[\psi\left(\frac{\alpha}{2}\right) - \psi(1) + \frac{1}{2} \ln \frac{l^2}{L^2} + \frac{2}{p^2 L^2} - \frac{3}{4} \right] \hat{h}_k^l(p) \right\} . \end{aligned} \quad (119)$$

From eqn. (119) we can read off the correlators

$$\langle \mathcal{O}(p) \rangle = \frac{\sqrt{6}\kappa l}{3L^2} \delta(p) , \quad (120)$$

$$\langle \mathcal{O}(p) \mathcal{O}(q) \rangle - \langle \mathcal{O}(p) \rangle \langle \mathcal{O}(q) \rangle = -\frac{\kappa}{l} \left[\psi\left(\frac{\alpha}{2}\right) - \psi(1) - \frac{4}{3p^2 L^2} \right] \delta(p+q) , \quad (121)$$

$$\langle T_j^i(-p) \mathcal{O}(p) \rangle = \frac{\sqrt{6}\kappa l}{3L^2} \left(\frac{2}{3p^2} P_j^i - \delta_j^i \right) , \quad (122)$$

$$\langle T_j^i(-p) T_l^k(p) \rangle = -\frac{\kappa l^3}{2} \Pi_{jl}^{ik} \left[\psi\left(\frac{\alpha}{2}\right) - \psi(1) - \frac{2}{3p^2 L^2} \right] , \quad (123)$$

where we have discarded some local terms. In particular, we have shifted the $p^2 = 0$ pole term proportional to $P_j^i P_l^k$ in the energy momentum tensor two-point function to the traceless transversal part. Notice that the term proportional to δ_j^i in $\langle T_j^i(-p) \mathcal{O}(p) \rangle$ has its origin in the source term inserted into the boundary field theory action functional. Thus, T_j^i is not exactly the

energy momentum operator of the field theory, and, when one considers the true operator, this unusual term does not appear [42]. Eqn. (121) has the opposite sign with respect to the result of DeWolfe and Freedman [28]. This is a result of adding the counter term (58), which contributes to the finite on-shell action. For large $|p|$, eqn. (121) approaches the conformal two-point function for operators of dimension $\Delta = 2$ (with the correct sign), so that we conclude that our sign is the correct one. The pole at $p^2 = 0$ can be interpreted by the existence of a massless dilaton from the broken conformal symmetry. The traceless transversal part of eqn. (123) has already been found in [11].

4 Conclusions

In this paper, we have presented a detailed calculation of one- and two-point correlation functions in holographic renormalization group flows dual to the Poincaré invariant background solutions of generic bulk theories containing scalars coupled to gravity. We were able to explicitly distinguish between operator and vev flows by means of the condensate in the boundary field theory. This result was achieved by carefully considering the counter terms, which must be added to the bulk field theory action in order to cancel divergences. For operators of dimension $\Delta > d/2$, we can describe both operator and vev flows, whereas for operators of dimension $\Delta = d/2$ we only have vev flows.

Our main improvement for two-point functions in holographic renormalization group flows is that we considered the sources of the metric and scalar fluctuations as independent boundary data, which couple as sources to the energy momentum tensor and scalar operators of the boundary field theory, respectively. Thus, our method results in the complete non-local correlation functions $\langle \mathcal{O}\mathcal{O} \rangle$, $\langle T_j^i \mathcal{O} \rangle$ and $\langle T_j^i T_l^k \rangle$. An analysis of these correlation functions yields important facts about the boundary field theory. For example, our analysis of the GPPZ flow in Sec. 3.1 confirmed the operator relation $T_i^i = \beta \mathcal{O}$, where β is the renormalized holographic beta function. Moreover, there should be a relation between the $P_j^i P_l^k$ part of $\langle T_j^i T_l^k \rangle$ and the c -function of the RG flow [46]. In order to obtain the correct contact terms in the correlators, one would have to carefully analyze *all* divergent terms in the on-shell action and to add necessary counter terms along the lines of [36]. This has been achieved very recently in [42].

We have chosen to apply our improved method to the GPPZ and Coulomb branch flows, which have already been studied in the literature, in order to be able to directly verify the known results and to improve them by finding the complete set of two-point functions. We leave the application of our method to other holographic renormalization group flows for the future. In particular, a flow that interpolates between UV and IR conformal fixed points would be

specially interesting, because the correlation functions should exhibit this flow in their dependence on the momentum p . In that respect, toy models, which do not necessarily contain the super potentials of true super gravity theories, might be very valuable. Furthermore, our method remains to be tested for a vev flow with $\Delta > d/2$.

Finally, we would like to mention several interesting questions. It remains an open problem how to describe holographically RG flows generated by operators of dimension Δ with $d/2 - 1 \leq \Delta < d/2$ ($d/2 - 1$ is the unitarity bound). For the AdS background, the method of using the Legendre transformed action as the generating functional [39,40] yields the correct conformal correlation functions, but it is not yet known whether it can be generalized to RG flows. One might expect that a background solution describing an operator flow with $d/2 < \Delta < d/2 + 1$ also describes a vev flow with a condensate of an operator of dimension $\Delta' = d - \Delta$. A technical problem is that the known examples of RG flows do not involve scalar operators with dimensions within this interval.

Another interesting question is to find “conjugate” flows and the relations of their field theoretical duals. By “conjugate” we mean the following. Whether a background solution describes an operator or a vev flow is determined by the super potential U , because of eqn. (14). Given a bulk potential V , the super potential is not unique, but there are several solutions. According to eqn. (6), one could change the overall sign of U , but this corresponds to simply changing the sign of r , so that $r \rightarrow -\infty$ becomes the asymptotically AdS region, and no new physics appears. It is more interesting to consider the expansion around the fixed point, eqn. (42). In general, there are two solutions for λ , $\lambda > d/2$ describing an operator flow, and $\lambda < d/2$ describing a vev flow. We call these two flows “conjugate,” and we conjecture that the corresponding boundary field theories are related to each other.

Acknowledgements

I am thankful to M. Bianchi, D.Z. Freedman and K. Skenderis for e-mail correspondence and their critical comments on the first version of this paper. In a very recent paper, they have considered the same problems and obtained results essentially identical with mine [42]. Their approach includes all counter terms, and thus yields the correct contact terms in the correlators, with which I was not very careful. Moreover, I would like to thank J. Erdmenger, A. Petkou, R. Pettorino and M. Porrati for useful discussions. Financial support by the European Commission RTN programme HPRN-CT-2000-00131, in which I am associated with INFN, Sezione di Frascati, is gratefully acknowledged.

A Geometric Relations for Hyper Surfaces

This appendix shall begin with a review of basic geometric relations for hyper surfaces [47]. A hyper surface in a space–time with coordinates X^μ ($\mu = 0, \dots, d$) is defined as a set of $d + 1$ functions, $X^\mu(x^i)$ ($i = 1, \dots, d$), where x^i are the coordinates of the hyper surface. Moreover, let $\tilde{g}_{\mu\nu}$ and g_{ij} be the metric tensors of space–time and the hyper surface, respectively. The tangents $X_i^\mu \equiv \partial_i X^\mu$ and the normal N^μ of the hyper surface satisfy the following orthogonality relations,

$$\tilde{g}_{\mu\nu} X_i^\mu X_j^\nu = g_{ij} , \quad (\text{A.1})$$

$$X_i^\mu N_\mu = 0 , \quad (\text{A.2})$$

$$N^\mu N_\mu = 1 . \quad (\text{A.3})$$

As in the main text, a tilde will be used to label quantities characterizing the $(d + 1)$ –dimensional space–time manifold, whereas those of the hyper surface remain unadorned.

The equations of Gauss and Weingarten define the second fundamental form, H_{ij} , of the hyper surface,

$$\partial_i X_j^\mu + \tilde{\Gamma}_{\lambda\nu}^\mu X_i^\lambda X_j^\nu - \Gamma_{ij}^k X_k^\mu = H_{ij} N^\mu , \quad (\text{A.4})$$

$$\partial_i N^\mu + \tilde{\Gamma}_{\lambda\nu}^\mu X_i^\lambda N^\nu = -H_i^j X_j^\mu . \quad (\text{A.5})$$

The second fundamental form describes the extrinsic curvature of the hyper surface and is related to the intrinsic curvature by another equation of Gauss,

$$\tilde{R}_{\mu\nu\lambda\rho} X_i^\mu X_j^\nu X_k^\lambda X_l^\rho = R_{ijkl} + H_{il} H_{jk} - H_{ik} H_{jl} . \quad (\text{A.6})$$

Furthermore, it satisfies the equation of Codazzi,

$$\tilde{R}_{\mu\nu\lambda\rho} X_i^\mu X_j^\nu N^\lambda X_k^\rho = \nabla_i H_{jk} - \nabla_j H_{ik} . \quad (\text{A.7})$$

The above formulae simplify, if (as in the familiar time slicing formalism [48,49]), we choose space–time coordinates such that

$$X^0 = r , \quad X^i = x^i . \quad (\text{A.8})$$

Then, the tangent vectors are given by $X_i^0 = 0$ and $X_i^j = \delta_i^j$. One conveniently splits up the space–time metric as (shown here for Euclidean signature)

$$\tilde{g}_{\mu\nu} = \begin{pmatrix} n_i n^i + n^2 & n_j \\ n_i & g_{ij} \end{pmatrix} , \quad (\text{A.9})$$

whose inverse is given by

$$\tilde{g}^{\mu\nu} = \frac{1}{n^2} \begin{pmatrix} 1 & -n^j \\ -n^i & n^2 g^{ij} + n^i n^j \end{pmatrix} . \quad (\text{A.10})$$

The matrix g^{ij} is the inverse of g_{ij} and is used to raise hyper surface indices. The quantities n and n^i are the lapse function and shift vector, respectively.

The normal vector N^μ satisfying the eqns. (A.2) and (A.3) is given by

$$N_\mu = (n, 0) , \quad N^\mu = \frac{1}{n}(1, -n^i) . \quad (\text{A.11})$$

Then, one obtains the second fundamental form from the equation of Gauss, eqn. (A.4), as

$$H_{ij} = -\frac{1}{2n} (\partial_r g_{ij} - \nabla_i n_j - \nabla_j n_i) . \quad (\text{A.12})$$

Finally, using the equations of Gauss and Weingarten, certain components of the space-time connections can be expressed as follows,

$$\tilde{\Gamma}_{ij}^r = \frac{1}{n} H_{ij} , \quad (\text{A.13})$$

$$\tilde{\Gamma}_{ij}^k = \Gamma_{ij}^k - \frac{n^k}{n} H_{ij} , \quad (\text{A.14})$$

$$\tilde{\Gamma}_{ir}^r = \frac{1}{n} \partial_i n + \frac{n^j}{n} H_{ij} , \quad (\text{A.15})$$

$$\tilde{\Gamma}_{ir}^k = \nabla_i n^k - \frac{n^k}{n} \partial_i n - n H_{ij} \left(g^{jk} + \frac{n^j n^k}{n^2} \right) . \quad (\text{A.16})$$

References

- [1] G. 't Hooft, [gr-qc/9310026](#).
- [2] L. Susskind, *J. Math. Phys.* **36** (1995) 6377, [hep-th/9409089](#).
- [3] J. M. Maldacena, *Adv. Theor. Math. Phys.* **2** (1998) 231, [hep-th/9711200](#).
- [4] S. S. Gubser, I. R. Klebanov and A. M. Polyakov, *Phys. Lett. B* **428** (1998) 105, [hep-th/9802109](#).
- [5] E. Witten, *Adv. Theor. Math. Phys.* **2** (1998) 253, [hep-th/9802150](#).
- [6] E. T. Akhmedov, *Phys. Lett. B* **442** (1998) 152, [hep-th/9806217](#).

- [7] L. Girardello, M. Petrini, M. Porrati and A. Zaffaroni, *J. High Energy Phys.* **12** (1998) 022, [hep-th/9810126](#).
- [8] A. Khavaev, K. Pilch and N. P. Warner, *Phys. Lett. B* **487** (2000) 14, [hep-th/9812035](#).
- [9] J. Distler and F. Zamora, *Adv. Theor. Math. Phys.* **2** (1999) 1405, [hep-th/9810206](#).
- [10] D. Z. Freedman, S. S. Gubser, K. Pilch and N. P. Warner, *Adv. Theor. Math. Phys.* **3** (1999) 363, [hep-th/9904017](#).
- [11] D. Z. Freedman, S. S. Gubser, K. Pilch and N. P. Warner, *J. High Energy Phys.* **07** (2000) 038, [hep-th/9906194](#).
- [12] L. Girardello, M. Petrini, M. Porrati and A. Zaffaroni, *Nucl. Phys. B* **569** (2000) 451, [hep-th/9909047](#).
- [13] J. Distler and F. Zamora, *J. High Energy Phys.* **05** (2000) 005, [hep-th/9911040](#).
- [14] E. Alvarez and C. Gómez, *Nucl. Phys. B* **541** (1999) 441, [hep-th/9807226](#).
- [15] M. Porrati and A. Starinets, *Phys. Lett. B* **454** (1999) 77, [hep-th/9903085](#).
- [16] B. Balasubrasmanian and P. Kraus, *Phys. Rev. Lett.* **83** (1999) 3605, [hep-th/9903190](#).
- [17] O. DeWolfe, D. Z. Freedman, S. S. Gubser and A. Karch, *Phys. Rev. D* **62** (2000) 046008, [hep-th/9909134](#).
- [18] V. Sahakian, *Phys. Rev. D* **62** (2000) 126011, [hep-th/9910099](#).
- [19] M. Petrini and A. Zaffaroni, [hep-th/0002172](#).
- [20] M. Porrati and A. Starinets, [hep-th/0009198](#).
- [21] M. Porrati and A. Starinets, *Phys. Lett. B* **498** (2001) 285, [hep-th/0009227](#).
- [22] J. Erdmenger, [hep-th/0103219](#).
- [23] A. Brandhuber and K. Sfetsos, *J. High Energy Phys.* **10** (1999) 013, [hep-th/9908116](#).
- [24] I. Chepelev and R. Roiban, *Phys. Lett. B* **462** (1999) 74, [hep-th/9906224](#).
- [25] S. B. Giddings and S. F. Ross, *Phys. Rev. D* **61** (2000) 024036, [hep-th/9907204](#).
- [26] R. C. Rashkov and K. S. Viswanathan, *Phys. Rev. D* **62** (2000) 046009, [hep-th/9911160](#).
- [27] D. Anselmi, L. Girardello, M. Porrati and A. Zaffaroni, *Phys. Lett. B* **481** (2000) 346, [hep-th/0002066](#).
- [28] O. DeWolfe and D. Z. Freedman, [hep-th/0002226](#).
- [29] G. Arutyunov, S. Frolov and S. Theisen, *Phys. Lett. B* **484** (2000) 295, [hep-th/0003116](#).
- [30] M. Bianchi, O. DeWolfe, D. Z. Freedman and K. Pilch, *J. High Energy Phys.* **01** (2001) 021, [hep-th/0009156](#).

- [31] A. Brandhuber and K. Sfetsos, *J. High Energy Phys.* **12**, (2000) 14, [hep-th/0010048](#).
- [32] K. Skenderis and P. K. Townsend, *Phys. Lett. B* **468** (1999) 46, [hep-th/9909070](#).
- [33] M. Henningson and K. Skenderis, *J. High Energy Phys.* **07** (1998) 023, [hep-th/9806087](#).
- [34] H. Liu and A. A. Tseytlin, *Nucl. Phys. B* **533** (1998) 88, [hep-th/9804083](#).
- [35] G. E. Arutyunov and S. A. Frolov, *Nucl. Phys. B* **544** (1999) 576, [hep-th/9806216](#).
- [36] S. de Haro, K. Skenderis and S. N. Solodukhin, *Commun. Math. Phys.* **217** (2000) 595, [hep-th/0002230](#).
- [37] J. de Boer, E. Verlinde and H. Verlinde, *J. High Energy Phys.* **08** (2000) 003, [hep-th/9912012](#).
- [38] P. Breitenlohner and D. Z. Freedman, *Ann. Phys. (N.Y.)* **144** (1982) 249.
- [39] I. R. Klebanov and E. Witten, *Nucl. Phys. B* **556** (1999) 89, [hep-th/9905104](#).
- [40] W. Mück and K. S. Viswanathan, *Phys. Rev. D* **60** (1999) 081901, [hep-th/9906155](#).
- [41] J. Kalkkinen, D. Martelli and W. Mück, *J. High Energy Phys.* **04** (2001) 036, [hep-th/0103111](#).
- [42] M. Bianchi, D. Z. Freedman and K. Skenderis, [hep-th/0105276](#).
- [43] M. Abramowitz and I. A. Stegun, eds., *Handbook of Mathematical Functions*, (Dover Publ., New York, 1965).
- [44] H. Osborn, *Nucl. Phys. B* **363** (1991) 486.
- [45] A. Brandhuber and K. Sfetsos, *Adv. Theor. Math. Phys.* **3** (1999) 851, [hep-th/9906201](#).
- [46] D. Anselmi, *Ann. Phys. (N.Y.)* **276** (1999) 361.
- [47] L. P. Eisenhart, *Riemannian Geometry*, (Princeton University Press, Princeton, 1964).
- [48] R. M. Wald, *General Relativity*, (University of Chicago Press, Chicago, 1984).
- [49] C. W. Misner, K. S. Thorne and J. A. Wheeler, *Gravitation*, (Freeman, San Francisco, 1973).